

Keep notes along the way

Putting pen to paper can help provide a chance for both personal reflection and a way to keep track of things along your loved one's recovery journey. Beginning to journal doesn't have to feel intimidating—it's all about what is most helpful for you.

On the following pages are two journaling templates to give you a place to start.

- **Self-Care Journal:** Record what you are doing, thinking, and feeling. It's intended to be used with the [Caregiver Self-Care Guide](#) to keep yourself healthy and balanced
- **My Loved One's Recovery Journal:** Keep track of events, successes, setbacks, and anything else you want to note along your loved one's recovery journey or discuss with their Treatment Team. It can be used alongside the [Your Episode Action Plan worksheet](#) to record any triggers, coping mechanisms, and early episode* warning signs. Mental health recovery is an ongoing process, not a single outcome. There are ups and downs along the way, but it is important to be hopeful

*An episode is defined as a relapse.



INDICATIONS

INVEGA HAFYERA® (6-month paliperidone palmitate) is a prescription medicine given by injection every 6 months by a healthcare professional and used to treat schizophrenia. INVEGA HAFYERA® is used in adults who have been treated with either:

- INVEGA SUSTENNA® (paliperidone palmitate) a 1-time-each-month paliperidone palmitate extended-release injectable suspension for at least 4 months
- INVEGA TRINZA® (paliperidone palmitate) a 1-time-every-3-months paliperidone palmitate extended-release injectable suspension for at least 3 months

INVEGA TRINZA® is a prescription medicine given by injection every 3 months by a healthcare professional and used to treat schizophrenia. INVEGA TRINZA® is used in people who have been adequately treated with INVEGA SUSTENNA® for at least 4 months.

INVEGA SUSTENNA® is a prescription medicine given by injection by a healthcare professional. INVEGA SUSTENNA® is used to treat schizophrenia in adults.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA®?

INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA® may cause serious side effects, including:

- **Increased risk of death in elderly people with dementia-related psychosis.** INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA® increase the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA® are not for the treatment of people with dementia-related psychosis.

Please read the additional Important Safety Information on pages 4-6 and the full Prescribing Information, including Boxed WARNING, for [INVEGA HAFYERA®](#), [INVEGA TRINZA®](#) and [INVEGA SUSTENNA®](#) and discuss any questions you have with your doctor.



Date: _____

What activities did I do today to better my physical well-being?

What did I do today that helped me feel emotionally stronger?

Who did I reach out to for help today, and why did I reach out?

What did I learn today? (Example: did you have any successes or setbacks today?
What can you take from those experiences?)

Did anything happen today with my loved one that affected how I feel about my role as a caregiver?

My Loved One's Recovery Journal



Date: _____

Medications Currently Taking: _____

What symptoms did my loved one experience today?

What triggers or stressors did my loved one encounter today?

What coping mechanisms did we use today? Were they helpful?

Did I see any episode warning signs today?

What progress or success—small or large—did my loved one make today towards their goals?

Other:

INDICATIONS

INVEGA HAFYERA® (6-month paliperidone palmitate) is a prescription medicine given by injection every 6 months by a healthcare professional and used to treat schizophrenia. INVEGA HAFYERA® is used in adults who have been treated with either:

- INVEGA SUSTENNA® (paliperidone palmitate) a 1-time-each-month paliperidone palmitate extended-release injectable suspension for at least 4 months
- INVEGA TRINZA® (paliperidone palmitate) a 1-time-every-3-months paliperidone palmitate extended-release injectable suspension for at least 3 months

INVEGA TRINZA® is a prescription medicine given by injection every 3 months by a healthcare professional and used to treat schizophrenia. INVEGA TRINZA® is used in people who have been adequately treated with INVEGA SUSTENNA® for at least 4 months.

INVEGA SUSTENNA® is a prescription medicine given by injection by a healthcare professional. INVEGA SUSTENNA® is used to treat schizophrenia in adults.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA®?

INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA® may cause serious side effects, including:

- **Increased risk of death in elderly people with dementia-related psychosis.** INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA® increase the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA® are not for the treatment of people with dementia-related psychosis.

Do not receive INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA® if you are allergic to paliperidone, paliperidone palmitate, risperidone, or any of the ingredients in INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA®. See the end of the Patient Information leaflet in the full Prescribing Information for a complete list of INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA® ingredients.

Before you receive INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA®, tell your healthcare professional about all your medical conditions, including if you:

- have had Neuroleptic Malignant Syndrome (NMS)
- have or have had heart problems, including a heart attack, heart failure, abnormal heart rhythm, or long QT syndrome
- have or have had low levels of potassium or magnesium in your blood
- have or have had uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia)
- have or have had kidney or liver problems
- have diabetes or have a family history of diabetes
- have Parkinson's disease or a type of dementia called Lewy Body Dementia
- have had a low white blood cell count
- have had problems with dizziness or fainting or are being treated for high blood pressure
- have or have had seizures or epilepsy
- have any other medical conditions
- are pregnant or plan to become pregnant. It is not known if INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA® will harm your unborn baby
 - If you become pregnant while taking INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA®, talk to your healthcare professional about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry>.

- Infants born to women who are treated with INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA® may experience symptoms such as tremors, irritability, excessive sleepiness, eye twitching, muscle spasms, decreased appetite, difficulty breathing, or abnormal movement of arms and legs. Let your healthcare professional know if these symptoms occur.
- are breastfeeding or plan to breastfeed. INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA® can pass into your breast milk. Talk to your healthcare professional about the best way to feed your baby if you receive INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA®.

Tell your healthcare professional about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA® may affect the way other medicines work, and other medicines may affect how INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA® works.

Your healthcare provider can tell you if it is safe to receive INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA® with your other medicines. Do not start or stop any medicines during treatment with INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA® without talking to your healthcare provider first. Know the medicines you take. Keep a list of them to show to your healthcare professional or pharmacist when you get a new medicine.

Patients (particularly the elderly) taking antipsychotics with certain health conditions or those on long-term therapy should be evaluated by their healthcare professional for the potential risk of falls.

How will I receive INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA®?

- Follow your treatment schedule exactly as your healthcare provider tells you to.
- Your healthcare provider will tell you how much you will receive and when you will receive it.

What should I avoid while receiving INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA®?

- INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA® may affect your ability to make decisions, think clearly, or react quickly. Do not drive, operate heavy machinery, or do other dangerous activities until you know how INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA® affects you.
- Avoid getting overheated or dehydrated.

INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA® may cause serious side effects, including:

- **See “What is the most important information I should know about INVEGA HAFYERA®, INVEGA TRINZA® and INVEGA SUSTENNA®?”**
- **stroke in elderly people (cerebrovascular problems) that can lead to death**
- **Neuroleptic Malignant Syndrome (NMS).** NMS is a rare but very serious problem that can happen in people who receive INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA®. NMS can cause death and must be treated in a hospital. Call your healthcare professional right away if you become severely ill and have any of these symptoms: high fever; severe muscle stiffness; confusion; loss of consciousness; changes in your breathing, heartbeat, and blood pressure.
- **problems with your heartbeat.** These heart problems can cause death. Call your healthcare professional right away if you have any of these symptoms: passing out or feeling like you will pass out, dizziness, or feeling as if your heart is pounding or missing beats.
- **uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia)**
- **metabolic changes.** Metabolic changes may include high blood sugar (hyperglycemia), diabetes mellitus and changes in the fat levels in your blood (dyslipidemia), and weight gain.
- **low blood pressure and fainting**
- **changes in your blood cell counts**
- **high level of prolactin in your blood (hyperprolactinemia).** INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA® may cause a rise in the blood levels of a hormone called prolactin (hyperprolactinemia) that may cause side effects including missed menstrual periods, leakage of milk from the breasts, development of breasts in men, or problems with erection.
- **problems thinking clearly and moving your body**
- **seizures**
- **difficulty swallowing that can cause food or liquid to get into your lungs**

- **prolonged or painful erection lasting more than 4 hours.** Call your healthcare professional or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours.
- **problems with control of your body temperature, especially when you exercise a lot or spend time doing things that make you warm. It is important for you to drink water to avoid dehydration.**

The most common side effects of INVEGA HAFYERA® include: injection site reactions, weight gain, headache, upper respiratory tract infections, feeling restlessness or difficulty sitting still, slow movements, tremors, stiffness and shuffling walk.

The most common side effects of INVEGA TRINZA® include: injection site reactions, weight gain, headache, upper respiratory tract infections, feeling restlessness or difficulty sitting still, slow movements, tremors, stiffness and shuffling walk.

The most common side effects of INVEGA SUSTENNA® include: injection site reactions; sleepiness or drowsiness; dizziness; feeling of inner restlessness or needing to be constantly moving; abnormal muscle movements, including tremor (shaking), shuffling, uncontrolled involuntary movements, and abnormal movements of your eyes.

Tell your healthcare professional if you have any side effect that bothers you or does not go away. These are not all the possible side effects of INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA®. For more information, ask your healthcare professional or pharmacist.

Call your healthcare professional for medical advice about side effects. You may report side effects of prescription drugs to the FDA at 1-800-FDA-1088.

General information about the safe and effective use of INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA®

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA® for a condition for which it was not prescribed. You can ask your pharmacist or healthcare professional for information about INVEGA HAFYERA®, INVEGA TRINZA® or INVEGA SUSTENNA® that is written for healthcare professionals.

For more information, go to www.invegahafyera.com, www.invegatrinza.com or www.invegasustenna.com or call 1-800-526-7736.

Please click to read the full Prescribing Information, including Boxed WARNING, for [INVEGA HAFYERA®](#), [INVEGA TRINZA®](#) and [INVEGA SUSTENNA®](#) and discuss any questions you have with your healthcare professional.

cp-256259v4